



The therapeutic role of proprioceptive training in neurological physiotherapy: An analytical study

Dr. Mohan Sharma¹, Dr. Jitendra Kumar², Dr Nidhi Sharma³, Dr. Arun Kumar Sharma⁴

¹ Assistant Professor, Star College of Physiotherapy, Jaipur, Rajasthan, India

² Assistant professor, Symbiosis College of Physiotherapy, Jaipur, Rajasthan, India

³ Lecturer, R R College of Physiotherapy, Jaipur, Rajasthan, India

⁴ Associate Professor, Symbiosis College of Physiotherapy Jaipur, Rajasthan, India

DOI: <https://doi.org/10.66856/ijyppe.2026.11.4.11106>

Abstract

Proprioception plays an important role in the control and coordination of human movement by providing information about joint position, body movement, and spatial orientation. Damage to the nervous system may disturb this sensory feedback and consequently affect balance, gait, coordination, postural stability, and functional independence. Conditions such as stroke, spinal cord injury, multiple sclerosis, traumatic brain injury, and other neurological disorders may therefore benefit from rehabilitation approaches that specifically address sensory-motor deficits. This analytical study explores the therapeutic significance of proprioceptive training in neurological physiotherapy, focusing on its potential contribution to postural control, balance, gait performance, coordination, and functional mobility. Different approaches, including weight-bearing exercises, joint-position activities, closed-chain exercises, balance and perturbation training, task-oriented practice, and technology-supported rehabilitation, are examined in relation to neurological recovery. The available evidence indicates that systematic proprioceptive training may improve the ability to receive, interpret, and respond to sensory information during movement and may consequently support better functional performance. Nevertheless, the effectiveness of training may differ according to the type and severity of neurological impairment, sensory deficits, cognitive abilities, exercise intensity, and patient participation. Proprioceptive training should therefore be adapted to individual functional needs and progressively incorporated into a broader rehabilitation programme. Combining proprioceptive exercises with conventional physiotherapy, strengthening, balance activities, and task-specific functional practice may provide a more comprehensive approach to neurological rehabilitation. Conclusively, proprioceptive training has considerable therapeutic relevance and can serve as an important component of patient-centred neurological physiotherapy.

Keywords: Proprioception, neurological physiotherapy, sensory-motor integration, balance, postural control

Introduction

Neurological disorders can substantially interfere with a person's ability to move, maintain balance, coordinate body movements, walk safely, and perform daily activities independently. Because these problems often persist beyond the acute stage of illness, physiotherapy and rehabilitation play an important role in improving functional ability and participation in everyday life. The World Health Organization (WHO) recognises rehabilitation as an essential component of care for people living with neurological conditions, including stroke, Parkinson's disease, spinal cord injury, traumatic brain injury, cerebral palsy, and other disorders affecting the nervous system [1]. Contemporary neurological physiotherapy therefore extends beyond improving muscle strength and joint mobility and also addresses the sensory mechanisms that contribute to accurate and controlled movement.

Proprioception is an important sensory component of human movement. It enables a person to recognise the position, direction, and movement of different parts of the body, even when visual information is limited. Sensory information arising from structures such as muscle spindles, Golgi tendon organs, joint receptors, and other mechanoreceptors is continuously processed to support posture, movement, and coordination. Neurological damage can disturb the transmission, processing, or integration of this information and may consequently affect the person's ability to control

body position and movement [2]. Clinically, this may be observed as poor postural stability, inaccurate limb positioning, abnormal gait patterns, and difficulty transferring weight, impaired coordination, and reduced ability to perform functional tasks without assistance.

Proprioceptive training aims to strengthen the relationship between sensory information and the motor responses required for controlled movement. In neurological physiotherapy, it can involve a range of activities, including weight-bearing exercises, joint-position awareness tasks, closed-kinetic-chain exercises, balance activities, rhythmic stabilisation, perturbation training, proprioceptive neuromuscular facilitation, and task-oriented movements [3]. The difficulty and complexity of these activities can be adjusted according to the patient's neurological condition, functional capacity, safety, and rehabilitation goals. For instance, a person recovering from stroke with weakness on one side may begin with supported weight transfer towards the affected limb. As control improves, treatment may progress to unsupported standing, reaching, stepping, directional changes, and gait-related tasks.

Research evidence indicates that proprioceptive interventions may provide clinically relevant benefits, particularly among individuals undergoing stroke rehabilitation. A meta-analysis involving 17 randomised controlled trials and 447 participants with stroke reported improvements in balance, walking speed, trunk control, and

basic functional mobility following proprioceptive training [4]. Further evidence has explored the value of combining proprioceptive approaches with other rehabilitation methods. A 2024 systematic review and network meta-analysis examined 64 randomised controlled trials involving 4,084 individuals with stroke-related lower-limb dysfunction. The findings indicated beneficial effects of different proprioceptive training combinations on balance, lower-limb motor performance, and walking ability [5]. These findings suggest that proprioceptive exercises may have greater practical value when incorporated into a broader, function-oriented rehabilitation programme rather than being applied as an independent treatment approach.

The contribution of proprioception becomes particularly important during balance and gait rehabilitation. Maintaining an upright posture requires the nervous system to continuously integrate information from the visual, vestibular, and somatosensory systems. When proprioceptive feedback is reduced or inaccurately interpreted, individuals may experience difficulty detecting changes in body position and adapting their movements to different environmental conditions [6]. Physiotherapists can address these limitations by progressively introducing activities that challenge weight distribution, postural reactions, limb loading, reaching, stepping, changes in the support surface, and dual-task performance. Such progression allows patients to develop better sensory-motor responses while gradually moving from controlled therapeutic activities towards functional movements required in everyday life. Evidence from systematic research on proprioceptive and dual-task interventions also suggests potential improvements in postural balance and gait and a possible reduction in fall-related risk among people recovering from stroke [7].

Appropriate assessment is necessary to determine whether proprioceptive rehabilitation is producing meaningful functional changes. In clinical practice, physiotherapists may evaluate joint-position awareness, kinaesthetic ability, postural control, balance, walking performance, and functional mobility. Standardised instruments such as the Berg Balance Scale, Timed Up and Go Test, 10-Metre Walk Test, 6-Minute Walk Test, Functional Reach Test, and Functional Ambulation Category can provide objective information about a patient's baseline status and progress during rehabilitation [8]. These measures are particularly useful because improvements in sensory-motor control should ultimately translate into practical abilities, such as standing independently, walking, and turning, negotiating stairs, changing direction, and moving safely within the community.

Despite its potential benefits, proprioceptive training cannot be considered a uniform intervention that produces identical results in every neurological patient. Individuals may differ considerably in the nature and severity of sensory impairment, muscle weakness, cognitive ability, balance capacity, fatigue, disease progression, and personal rehabilitation goals. Consequently, physiotherapists need to individualise the selection, intensity, repetition, duration, sensory challenge, external feedback, and progression of proprioceptive exercises. Another important issue is the methodological variation observed across existing studies.

Differences in intervention protocols, patient characteristics, treatment duration, and outcome measures can make comparisons between studies difficult. Although previous research has reported positive changes in sensorimotor and functional outcomes, further investigation is required to determine which proprioceptive strategies are most effective for specific neurological conditions and patient populations. In view of these considerations, the present analytical study examines the therapeutic contribution of proprioceptive training within neurological physiotherapy. Particular emphasis is placed on its potential effects on balance, gait, postural control, motor coordination, functional mobility, and independence in daily activities. The study also considers commonly used training techniques, clinical assessment methods, available research evidence, practical challenges, and existing gaps in the literature. A clearer understanding of these aspects can support the appropriate integration of proprioceptive principles into individualised neurological rehabilitation and help physiotherapists combine sensory-motor interventions with conventional and task-specific therapeutic approaches.

Proprioceptive Training in Neurological Physiotherapy

Proprioceptive training is used in neurological physiotherapy to improve the ability to recognise body position, movement, and changes in posture and to use this information for better motor control. Neurological disorders may disturb sensory feedback and its integration with voluntary movement, leading to problems with balance, gait, coordination, and functional mobility. Proprioceptive exercises therefore provide controlled sensory and movement experiences that can be incorporated into a broader rehabilitation programme. The aim is not only to improve sensory awareness but also to help patients use this information during meaningful functional activities.

Principles and Mechanisms of Proprioceptive Training

An effective proprioceptive programme should be progressive, repetitive, functional, and suited to the patient's abilities. Training may begin with simple movements performed with support and gradually advance towards more challenging activities. The therapist can increase the difficulty by reducing external support, changing the base of support, altering visual input, increasing movement complexity, or introducing controlled disturbances. The underlying goal is to improve communication between sensory information and motor responses. Repeated practice may help patients develop more accurate movement strategies and improve their ability to respond to changes in body position [9]. Proprioceptive training should therefore be viewed as part of motor learning and functional rehabilitation rather than as an isolated exercise technique.

Major Proprioceptive Training Approaches

Several techniques can be used according to the patient's neurological impairment and functional needs. Weight-bearing and closed-chain exercises such as supported standing, weight shifting, sit-to-stand practice, mini-squats, and stepping provide sensory input through the muscles and joints while encouraging controlled limb loading. However,

Balance and postural exercises may include tandem standing, reaching, directional stepping, turning, and controlled changes in the support surface ^[10]. These activities challenge the patient's ability to maintain stability while processing sensory information.

Rhythmic stabilisation and proprioceptive neuromuscular facilitation (PNF) may be used to improve postural control, coordinated muscle activity, and movement patterns. Perturbation training introduces carefully controlled changes in balance so that patients can practise appropriate postural reactions ^[11]. The choice of technique should depend on the patient's condition, current level of function, safety, and rehabilitation goals rather than following one fixed protocol.

Task-Specific and Functional Proprioceptive Training

Proprioceptive training becomes more clinically meaningful when it is connected with activities that patients need to perform in daily life. Exercises can therefore be incorporated into sit-to-stand transfers, walking, stair negotiation, reaching, turning, stepping in different directions, and changes in walking speed. For example, a person recovering from stroke may initially practise controlled weight transfer towards the weaker side ^[12]. This can progress to stepping, reaching while standing, turning, and walking. Similarly, a person with Parkinson's disease may practise postural control, stepping, turning, and movement initiation within functional tasks. Such task-specific practice allows sensory information to be used during movements that are relevant to everyday independence ^[13].

Application across Neurological Conditions

The application of proprioceptive training varies according to the neurological condition. In stroke, treatment commonly focuses on weight distribution, balance, limb awareness, gait, and functional mobility. In Parkinson's disease, proprioceptive and postural activities may be combined with stepping, turning, balance, and gait exercises. For individuals with spinal cord injury, appropriate sensory-motor activities may include supported standing, weight shifting, transfers, and gait-related training according to the level and completeness of injury. In multiple sclerosis, training may

need to be adjusted according to fatigue, balance impairment, weakness, and sensory changes. Patients with traumatic brain injury may benefit from activities addressing balance, coordination, body awareness, and functional movement ^[14].

Technology-Assisted Proprioceptive Rehabilitation

Technology can provide additional opportunities for sensory-motor training. Virtual reality, interactive balance systems, wearable sensors, robotic devices, and motion-feedback systems can provide information about body position, movement, balance, and performance ^[15]. These approaches may increase repetition, provide immediate feedback, and make exercises more engaging. However, technology should support rather than replace clinical physiotherapy. The physiotherapist remains responsible for selecting suitable activities, adjusting difficulty, monitoring safety, and interpreting changes in functional performance. Factors such as cost, equipment availability, patient cognition, accessibility, and clinical setting should also be considered ^[16].

Individualisation, Progression and Safety

Proprioceptive rehabilitation should be individually planned. Factors such as neurological diagnosis, sensory impairment, muscle strength, balance ability, cognition, fatigue, fear of falling, and functional goals can influence treatment selection. Progression may involve increasing repetitions, reducing physical support, changing the base of support, reducing visual dependence, increasing movement complexity, or introducing dual-task activities. Exercises should be sufficiently challenging to promote adaptation but should not create unnecessary risk. Close supervision is particularly important for patients with significant balance impairment or a history of falls ^[17].

Measurable Outcome Assessment

Objective assessment is important for determining whether proprioceptive training produces meaningful changes. Measurements should ideally be obtained before treatment, after the intervention period, and, where possible, during follow-up ^[18].

Outcome Area	Assessment Tool	What It Measures
Proprioceptive awareness	Joint-position error assessment	Accuracy of detecting joint position
Functional balance	Berg Balance Scale (BBS)	Ability to perform functional balance tasks
Dynamic mobility	Timed Up and Go (TUG)	Functional mobility and dynamic balance
Walking speed	10-Metre Walk Test (10MWT)	Gait speed
Walking endurance	6-Minute Walk Test (6MWT)	Walking capacity and endurance
Functional reach	Functional Reach Test	Limits of forward stability
Walking independence	Functional Ambulation Category (FAC)	Level of assistance required for walking
Motor recovery	Fugl-Meyer Assessment	Motor and sensory impairment, particularly after stroke
Functional independence	Appropriate ADL/independence scale	Ability to perform daily activities
Falls	Fall-frequency recording	Number of falls during a defined period

These measures help convert changes in balance, gait, motor control, and mobility into observable and measurable outcomes. Where possible, objective scores should be combined with functional observation and patient-reported changes to provide a broader picture of rehabilitation progress.

Clinical Integration

Proprioceptive training is most useful when integrated with other components of neurological physiotherapy rather than being used in isolation. Strengthening, range-of-motion exercises, balance training, gait rehabilitation, functional task practice, and patient education can be combined according to individual needs.

Outcome Measures and Measurement Properties ^[19]

Tool	Domain	Reliability	Validity	Responsiveness	Clinical Use
Berg Balance Scale (BBS)	Balance	High	Good	Good	Balance and postural control
Timed Up and Go (TUG)	Mobility	Excellent	Good	Good	Transfers and dynamic mobility
10-Meter Walk Test (10MWT)	Gait speed	High	Good	Good	Walking speed
6-Minute Walk Test (6MWT)	Endurance	High	Good	Good	Walking capacity
Functional Reach Test (FRT)	Stability	Moderate–High	Good	Moderate	Limits of stability
Functional Ambulation Category (FAC)	Walking independence	Excellent	Good	Good	Walking assistance
Fugl-Meyer Assessment (FMA)	Motor recovery	Excellent	Strong	Good	Motor and sensory recovery
Joint-Position Error Assessment	Proprioception	Variable	Direct	Potentially good	Proprioceptive accuracy

Note: Reliability, responsiveness, MDC (Minimal Detectable Change), and MCID (Minimal Clinically Important Difference) may vary according to neurological condition, disease stage, testing procedure, and patient characteristics.

Clinical Implications, Challenges and Research Gaps

Proprioceptive training can play an important role in neurological physiotherapy because problems with body-position awareness and sensory-motor control may affect balance, walking, coordination, and independence. However, the benefit of training depends on proper assessment, suitable exercise selection, safe progression, and its combination with other rehabilitation activities. The programme should therefore be adapted to each patient's neurological condition, sensory and motor abilities, balance, fatigue, cognitive status, and treatment goals.

Clinical Implications

Proprioceptive exercises are more useful when they are combined with strengthening, balance exercises, gait training, and functional activities. Simple activities such as weight shifting, standing, stepping, reaching, transfers, turning, and walking can help patients use sensory information during movement. The main aim should be to improve practical movement and daily function rather than focusing only on proprioceptive ability.

Regular assessment is important for deciding whether treatment should be continued, progressed, or modified. The Berg Balance Scale, Timed Up and Go Test, 10-Meter Walk Test, 6-Minute Walk Test, and Functional Ambulation Category can be used to monitor changes in balance, mobility, walking speed, endurance, and walking independence. Improvement should be considered not only in terms of test scores but also in terms of whether the patient can perform everyday activities more safely and independently.

Challenges and Clinical Limitations

Patients with neurological disorders do not have the same rehabilitation needs. Some may have sensory loss, weakness, spasticity, fatigue, poor coordination, cognitive difficulties, or fear of falling. Because of these differences, one proprioceptive training programme cannot be expected to work equally well for everyone. Exercises that are too difficult may increase instability and safety risks, while exercises that are too easy may provide little therapeutic challenge. Careful supervision and gradual progression are therefore important.

There are also practical difficulties in providing proprioceptive training. Limited therapy time, lack of specialised equipment, differences in therapist experience, poor patient adherence, and limited resources may affect treatment. Technologies such as virtual reality, wearable sensors, and robotic devices can provide additional feedback

and repetition, but their use may be restricted by cost, availability, technical requirements, and the patient's ability to use them safely.

Research Gaps

Although proprioceptive training has been studied in neurological rehabilitation, the available research is not fully consistent. Studies use different types of exercises, treatment durations, training frequencies, intensities, and outcome measures. In many studies, proprioceptive exercises are combined with balance training, strengthening, gait training, or conventional physiotherapy. As a result, it can be difficult to determine how much of the improvement is specifically related to proprioceptive training.

Many studies also include small numbers of participants and provide limited follow-up after treatment. More research is needed to identify the most suitable training dose and to determine whether improvements remain after the intervention ends. Future studies should also examine practical outcomes such as falls, community walking, activities of daily living, participation, and long-term independence.

Future Research Directions

Future research should develop clear and reproducible proprioceptive training programmes for different neurological conditions. Larger clinical trials with suitable control groups and longer follow-up periods are needed. Researchers should compare different training intensities and durations and use reliable, valid, and responsive outcome measures.

Further studies should also explore whether technology-assisted proprioceptive training provides greater benefits than conventional approaches. Importantly, research should determine whether improvements in proprioceptive ability lead to better balance, walking, daily activities, and long-term functional independence.

Conclusion

Proprioceptive training is an important part of neurological physiotherapy because it can help patients improve body awareness, movement control, balance, coordination, and mobility. Neurological disorders may affect a person's ability to understand the position and movement of different parts of the body. This can make walking, maintaining balance, and performing daily activities more difficult. Proprioceptive exercises can help reduce these difficulties when they are selected according to the patient's condition, abilities, and rehabilitation needs.

Proprioceptive training works best when it is combined with other physiotherapy approaches such as strengthening, balance exercises, gait training, and functional activities. This combination helps patients use improved sensory and movement control during everyday tasks. Tools such as the Berg Balance Scale, Timed Up and Go Test, 10-Meter Walk Test, 6-Minute Walk Test, and Functional Ambulation Category can be used to monitor changes in balance, mobility, walking, and independence.

However, current research has some limitations. Studies use different exercises, treatment durations, training intensities, and assessment methods, making their findings difficult to compare. Many studies also have small sample sizes and limited long-term follow-up. More research is therefore needed to identify the most suitable training methods and determine whether the benefits continue after treatment.

Conclusively, proprioceptive training should be used as part of a complete neurological rehabilitation programme rather than as a separate treatment. When exercises are planned according to individual needs, progressed safely, and linked with daily functional activities, proprioceptive training can help improve balance, mobility, independence, and participation in everyday life.

References

- World Health Organization. Package of Interventions for Rehabilitation: Module 3: Neurological Conditions. Geneva: World Health Organization,2023. <https://www.who.int/publications/i/item/9789240071131>.
- Proske U, Gandevia SC. The Proprioceptive Senses: Their Roles in Signalling Body Shape, Body Position and Movement, and Muscle Force. *Physiological Reviews*,2012;92(4):1651–1697. <https://doi.org/10.1152/physrev.00048.2011>.
- Aman CM, Elangovan RA, Stuart V, *et al.* The Effectiveness of Proprioceptive Training for Improving Motor Function: A Systematic Review. *Frontiers in Human Neuroscience*,2014;8:1075. <https://doi.org/10.3389/fnhum.2014.01075>.
- Apriliyasari RW, Truong PV, Tsai PS. Effects of Proprioceptive Training for People with Stroke: A Meta-Analysis of Randomized Controlled Trials. *Clinical Rehabilitation*,2022;36(4):431–448. <https://doi.org/10.1177/02692155211057656>.
- Zheng K, Li L, Zhou Y, Gong X, Zheng G, Guo L. Optimal Proprioceptive Training Combined with Rehabilitation Regimen for Lower Limb Dysfunction in Stroke Patients: A Systematic Review and Network Meta-Analysis. *Frontiers in Neurology*,2024;15:1503585. <https://doi.org/10.3389/fneur.2024.1503585>.
- Frasson F, Pizzigalli A, *et al.* Postural Control and Proprioception in Healthy Adults: A Systematic Review. *Journal of Clinical Medicine*,2023;12(6):2300. <https://doi.org/10.3390/jcm12062300>.
- Chiaromonte R, Bonfiglio M, Leonforte P, Coltraro GL, Guerrero CS, Vecchio M. Proprioceptive and Dual-Task Training: The Key of Stroke Rehabilitation, A Systematic Review. *Journal of Functional Morphology and Kinesiology*,2022;7(3):53. <https://doi.org/10.3390/jfmk7030053>.
- Park S, *et al.* Assessment of Lower Limb Motor Function, Ambulation, and Balance after Stroke. *Brain & Neurorehabilitation*,2022;15(2):e20. <https://doi.org/10.12786/bn.2022.15.e20>.
- Winter L, Huang Y, Sertic M, Konczak J. The Effectiveness of Proprioceptive Training for Improving Motor Performance and Motor Dysfunction: A Systematic Review. *Frontiers in Rehabilitation Sciences*,2022;3:830166. <https://doi.org/10.3389/fresc.2022.830166>.
- Ada L, Dean CM, Lindley E. Randomized Trial of Proprioceptive Training for Stroke Patients. *Clinical Rehabilitation*,2011;25(8):709–718. <https://doi.org/10.1177/0269215510396048>.
- Nguyen PT, Chou LW, Hsieh YL. Proprioceptive Neuromuscular Facilitation-Based Physical Therapy on the Improvement of Balance and Gait in Patients with Chronic Stroke: A Systematic Review and Meta-Analysis. *Life*,2022;12(6):882. <https://doi.org/10.3390/life12060882>.
- Lee MH, Lee DY. Effects of Task-Oriented Training on Gait Outcomes and Balance in Individuals with Stroke: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Journal of Clinical Medicine*,2025;14(24):8766. DOI: 10.3390/jcm14248766
- Perry SIB, Nelissen PM, Siemonsma P, Lucas C. The Effect of Functional-Task Training on Activities of Daily Living for People with Parkinson's Disease: A Systematic Review with Meta-Analysis. *Complementary Therapies in Medicine*,2019;42:312–321. DOI 10.1016/j.ctim.2018.12.008.
- Mason HK, *et al.* Exercise Prescription for Patients with Multiple Sclerosis; Potential Benefits and Practical Recommendations. *BMC Neurology*,2017;17:185. DOI: 10.1186/s12883-017-0960-0
- Laver KE, *et al.* Virtual Reality for Stroke Rehabilitation. *Cochrane Database of Systematic Reviews*,2025,6:CD008349. DOI 10.1002/14651858.CD008349.pub5.
- Shen J, *et al.* Effects of Virtual Reality–Based Exercise on Balance in Patients with Stroke: A Systematic Review and Meta-analysis. *American Journal of Physical Medicine & Rehabilitation*,2023;102(4):316–322. DOI 10.1097/PHM.0000000000002096.
- Winter L, Huang Q, Sertic JVL, Konczak J. The Effectiveness of Proprioceptive Training for Improving Motor Performance and Motor Dysfunction: A Systematic Review. *Frontiers in Rehabilitation Sciences*,2022;3:830166. DOI 10.3389/fresc.2022.830166.
- Van Crielinge T, *et al.* Standardized Measurement of Balance and Mobility Post-Stroke: Consensus-Based Core Recommendations from the Third Stroke Recovery and Rehabilitation Roundtable. *Neurorehabilitation and Neural Repair*,2024;38(1):41–51. DOI 10.1177/15459683231209154.
- Hillier HM, Holle AJ, Walsh RSC. Assessing Proprioception: A Systematic Review of Possibilities. *Journal of Rehabilitation Medicine*,2015;47(10):905–918. DOI 10.2340/16501977-1976.